

The Pion Light-Cone Wave Function Φ_π on the lattice: a partonic signal?

A. Abada^a, Ph. Boucaud^a, G. Herdoiza^a, J.P. Leroy^a, J. Micheli^a,
O. Pène^a and J. Rodríguez-Quintero^{b *}

^aLaboratoire de Physique Théorique[†]
Université Paris-Sud, Bâtiment 210, 91405 Orsay Cedex, France

^bDpto. de Física Aplicada e Ingeniería eléctrica
E.P.S. La Rábida, Universidad de Huelva, 21819 Palos de la fra., Spain

We determine the conditions required to study the pion light-cone wave function Φ_π with a new method: a direct display of the partons constituting the pion. We present the preliminary results of a lattice computation of Φ_π following this direction. An auxiliary scalar-quark is introduced. The spectroscopy of its bound states is studied. We observe some indications of a partonic behavior of the system of this scalar-quark and the anti-quark.

INTRODUCTION

The light-cone wave function (LCWF), $\Phi_\pi(u)$, plays an important role in the study [1] of pions with large momenta P_π . It is related to the probability of finding in the pion a quark and an anti-quark moving in the same direction than the pion with momenta uP_π and $(1-u)P_\pi$ respectively. Within this picture, the two valence quarks are frozen in a transverse region of size Λ_{QCD} . To extract the partonic information contained in $\Phi_\pi(u)$ a “kick” is applied to the pion with a large transfer momentum $\vec{q}$. This is for example what has been done in the E791 experiment [2] for the diffractive dissociation of a pion in two jets.

The LCWF Φ_π depends on a momentum scale which is typically P_π . In the large pion momentum frame, Φ_π describes the leading term of the $\Lambda_{\text{QCD}}^2/P_\pi^2$ expansion of the full pion wave function. It contributes only to the valence configuration $|q\bar{q}\rangle$ of the Fock space expansion. Φ_π contains all the terms of the $1/\log(P_\pi^2/\Lambda_{\text{QCD}}^2)$ expansion and is defined by an expression *à la* Bethe-Salpeter involving the pion to vacuum matrix element of a non-local operator

$$\langle 0 | \bar{d}(0) \mathcal{P} \left[\exp(i \int_x^0 d\tau_\mu A^\mu) \right] \gamma_\mu \gamma_5 u(x) | \pi(P_\pi) \rangle_{x^2=0}$$

*Talk given by Gregorio Herdoiza.

[†]Unité Mixte de Recherche du CNRS - UMR 8627.

$$= -i P_{\pi\mu} f_\pi \int_0^1 du e^{-iuP_\pi \cdot x} \Phi_\pi(u) \quad (1)$$

The Wilson string in the square bracket ensures the gauge invariance of the matrix element. Φ_π is a non-perturbative function, normalized by the condition $\int du \Phi_\pi(u) = 1$.

An important particular case is given by keeping only the dominant term in the expansion of $\mathcal{O}(1/\log(P_\pi^2/\Lambda_{\text{QCD}}^2))$. It leads to the asymptotic form of Φ_π , $\Phi_\pi^{\text{as}}(u) = 6u(1-u)$, which is fully computed by perturbative QCD¹.

On the lattice, the first method to study the LCWF is to examine the moments $\mathcal{M}_n$ of Φ_π [3], $\mathcal{M}_n = \int_0^1 du u^n \Phi_\pi(u)$. This requires the computation of the matrix elements of local operators. For high n , $\mathcal{M}_n$ involves operators with high order derivatives, which are increasingly difficult to discretize and renormalize. It was therefore proposed in [4] to calculate Φ_π by studying directly the *partonic* constituents of the pion. Following this idea, we have begun our work by stating the conditions needed to extract Φ_π , and then achieved a first and preliminary attempt to isolate a partonic signal on the lattice [5].

1. FEASIBILITY CONDITIONS

The r.h.s of eq. (1) is computed *via* a three-point Green function (see eq. (2) and fig. 1). The

¹ f_π being the only non-perturbative quantity on this limit.

Wilson string needed to insure gauge invariance is replaced by a *scalar coloured propagator* $S(\vec{x}_t, t; 0)$ with the colour quantum numbers of a quark, but this choice is not unique.




Figure 1. Three-point Green function computed on the lattice.

The active quark is transformed into the scalar-quark at time t and receives a large momentum transfer $\vec{q}$.

Exhibiting a partonic behavior on the lattice and thus computing directly Φ_π requires several conditions which have been derived in [5] and are listed below without their derivation.

First condition: large pion momentum, $P_\pi \gg \Lambda_{\text{QCD}}$, to have a picture of a “frozen” pion with a pair of almost colinear valence quarks. This means that we will have to deal with two scales, Λ_{QCD} and P_π , on the lattice.

Second condition: the scalar-quark has to be an energetic parton, $E_s \gg \Lambda_{\text{QCD}}$.

Third condition: the time t must be small to prevent the loss of quantum coherence between partons and the hadronisation. In particular we need $t \ll (1/\Lambda_{\text{QCD}}) \times (E_s/\Lambda_{\text{QCD}})$: in the pion rest frame, the partonic behavior lives at a time $\tau \ll 1/\Lambda_{\text{QCD}}$; the factor E_s/Λ_{QCD} acts as a sort of Lorentz factor. When t becomes large, the partonic system is replaced by a hadronic state, a bound state made of a quark and a scalar-quark that we will write $\tilde{\pi}$.

Last condition stemming from finite volume restrictions on the lattice: in the continuum the momentum fraction $u \in [0, 1]$, but due to momentum discretization on the lattice, $P_\pi = 2\pi/L_x \times n_\pi$, we will only cover a discrete set of values of u . Discarding $u = 0$ or 1 , because partons at rest is meaningless, we deal with $n_\pi \geq 2$. If $n_\pi = 2$, we are left with the only value $u = 1/2$.

With all these conditions verified, we compute the three-point Green function F^μ :

$$F^\mu(\vec{p}_\pi, \vec{q}; t) \equiv \int d^3x_\pi d^3x_t e^{-i\vec{p}_\pi \cdot \vec{x}_\pi} e^{-i\vec{q} \cdot \vec{x}_t} e^{E_\pi(t_\pi - t)} \times \langle 0 | P_5(\vec{x}_\pi, t_\pi) u(\vec{x}_t, t) S(\vec{x}_t, t; 0) \gamma_\mu \gamma_5 \bar{d}(0) | 0 \rangle \propto p_{\pi\mu} f_\pi \sum_{u_i} \frac{e^{-(E_s + (1-u_i)E_\pi)t}}{2E_s(u_i)} \Phi_\pi(u_i) \quad (2)$$

The matrix element now contains the interpolating field of the pion P_5 at the fixed time t_π and the scalar propagator. We have also included the inverse pion propagator $e^{E_\pi(t_\pi - t)}$ and required the time interval $t_\pi - t$ to be large in order to have, at the small times t , an on shell pion. In the r.h.s of eq. (2) we have replaced the integral over u by the sum because of the discretization constraint. It is important to note that the fraction which multiplies Φ_π can be understood within a *partonic picture* as the product of the inverse scalar propagator $e^{-E_s t}/2E_s$ times the inverse anti-quark propagator $e^{-(1-u_i)E_\pi t}$.

On the lattice we have tried to fulfill as much as possible the restrictive conditions just presented. This seriously constrain the simulation and turn the search of a signal into a delicate task.

2. LATTICE RESULTS

The simulation was performed on a $16^3 \times 40$ lattice ($\beta = 6.0$, 100 configurations) in the quenched approximation. We selected light parton masses ($\kappa_q \in \{0.1333, 0.1339\}$ for the quarks and $\kappa_s \in \{0.1428, 0.1430, 0.1431\}$ for the scalar) and the following values for $\vec{P}_\pi$, $\vec{P}_\pi \in \{(0, 0, 0); (1, 0, 0); (1, 1, 0); (2, 0, 0)\}$ in units of $2\pi/L_x$. A large set of transfer momenta $\vec{q}$ was chosen.

As we have introduced in eq. (2) a non physical object, the scalar-quark, the first step of the analysis was to test its hadronic behavior. We measured therefore the two-point Green functions for the $\tilde{\pi}$ and the scalar-scalar states. These two-point functions show the characteristic exponential signal of hadrons and verify the Einstein spectral law. Interestingly, the spectroscopy of the scalar reveals that these non-conventional bound states behave as real hadrons.

We will now consider the three-point Green




Figure 2. Slopes of the partonic ratio R_p and of the hadronic ratio R_h in the small time t interval $[0, 4]$. The horizontal axis displays the set of momenta $\vec{q}$ labelled with an arbitrary index.

function $F^0(\vec{P}_\pi, \vec{q}; t)$, eq. (2). As already justified, we use the momentum $\vec{P}_\pi = (2, 0, 0) \times 2\pi/L_x$ in order to have $u = 1/2$ although large momenta bring up some noise in a lattice signal.

We will try to give a partonic interpretation of the data in the short time t region by building the following ratios

$$R_p = \frac{F^0(\vec{p}_\pi, \vec{q}; t)}{\left[p_\pi^0 f_\pi \frac{e^{-(E_\pi/2 + E_s)t}}{2E_s} \Phi_\pi(1/2) \right]}$$

$$R_h = \frac{F^0(\vec{p}_\pi, \vec{q}; t)}{[e^{-E_\pi t}]}$$

We will look for plateaus in time of these quantities for the whole set of momenta $\vec{q}$. A plateau for R_p (R_h) would be a sign of a partonic (hadronic) behavior of the antiquark–scalar system. In fig. 2, we plot the *slopes* of the ratios R_p and R_h in the short time region $t \in [0, 4]$, for our set of momenta $\vec{q}$. A value of the slope close to zero suggests a plateau in time. By comparing both plots, we observe that the partonic slopes are closer to zero than the hadronic ones. According to our data, the partonic interpretation is somehow favoured: $\chi^2/\text{d.o.f} = 1.9$ for a vanishing slope compared to 4.1 in the hadronic case. Nevertheless, we are clearly not yet in a position to give a fully convincing claim.

As a cross-check of the previous result, we have examined the product $F^0(\vec{p}_\pi = (2, 0, 0), \vec{q}; t = 0) \times E_s(1/2)$. Within the partonic picture, we expect this product to be independent of the transfer momentum $\vec{q}$ (see eq. (2)). This is indeed what we find when we plot in fig. 3 this product for our full set of momenta $\vec{q}$. The data show the expected constancy around the average represented by the solid horizontal line.





Figure 3. Values of $F^0(\vec{p}_\pi = (2, 0, 0), \vec{q}; t = 0) \times E_s(1/2)$ for our full set of momenta $\vec{q}$.

CONCLUSION

We wanted to check the feasibility on the lattice of a direct method to determine non-perturbative objects like the pion light-cone wave function Φ_π . We have stated the conditions for this study and we have seen that they induce restrictive constraints and hierarchies which turn the search of a clear lattice signal into a difficult task. We have found an interesting spectroscopy of bound states involving a scalar-quark, and we have a hint of a partonic behavior of the system of the scalar and the anti-quark by considering the three-point Green function related to Φ_π . This signal will continue to be tested in a further analysis. Work supported by the European Community's Human potential programme under HPRN-CT-2000-00145 Hadrons/LatticeQCD.

REFERENCES

1. S. J. Brodsky *et al.*, Phys.Lett.**B91** (1980) 239.
2. E. M. Aitala *et al.* [E791 Collaboration], Phys.Rev.Lett.**86** (2001) 4768.
3. G. Martinelli *et al.*, Nucl.Phys.**B316** (1989) 355.
4. U. Aglietti *et al.*, Phys.Lett.**B441** (1998) 371.
5. A. Abada *et al.*, Phys.Rev.**D64** (2001) 074511.